PORTLAND — MULTNOMAH COUNTY





Adult Men: Weighted Estimates from a Probability Sample of Booked Arrestees 2003 Adult Male Program Findings

Facilities Sampled: 1 **Total Facilities:**

Total Number Booked in Sampled Facilities During Data Collection Period: 2703

Completed Interviews in Sample: Interview Refusal Rate: 14.5% 564 Completed Urine Tests in Sample: Urine Refusal Rate: 527 3.4%

1. Age					
	<21	21–25	26-30	31–35	36+
	11.7%	19.7%	15.2%	18.6%	34.9%

2. Race/Ethnicity									
Black	White	Hispanic	Other						
20.4%	57.5%	10.9%	11.1%						

	Tota	Sample		Age				Race/Ethnicity			
	%	95% CI	<21	21-25	26-30	31-35	36+	Black	White	Hispanic	Other
Alcohol	7.0	(+/-2)	4.9	2.8	6.7	7.9	9.6	3.1	9.1	2.1	7.8
Cocaine	29.7	(+/-4)	10.7	21.7	31.1	31.7	38.8	53.8	21.7	28.4	27.8
Marijuana	38.4	(+/-4)	61.2	52.1	46.0	29.8	24.8	46.3	35.4	35.5	42.5
Opiates	15.0	(+/-3)	10.2	16.0	17.0	15.8	14.7	5.5	18.1	10.4	21.2
Methamphetamine	25.4	(+/-4)	20.0	25.8	32.0	34.1	19.4	4.9	35.1	15.4	22.7
PCP	0.0	(+/-0)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Any Drug ^b	74.7	(+/-4)	78.9	75.6	81.5	78.2	68.2	80.0	75.1	63.7	73.9
Any Drug (excluding alcohol)	71.5	(+/-4)	76.4	74.1	79.2	74.1	63.9	79.2	70.7	61.6	71.1
Multiple Drugs ^c	32.5	(+/-4)	27.0	33.1	34.6	33.7	32.4	31.5	34.5	20.4	35.6
Multiple Drugs (excluding alcohol)	29.8	(+/-4)	25.7	32.6	33.8	31.1	27.1	29.1	31.1	20.4	33.3

	Walant	Dramartis	Device	Domestic	Driving While	
	Violent	Property	Drug	Violence	Intoxicated	Other
Alcohol	8.1	5.6	9.5	11.2	15.8	5.9
Cocaine	17.7	28.6	41.9	18.5	35.1	35.2
Marijuana	37.0	34.9	38.8	36.9	49.2	39.7
Opiates	8.6	21.0	15.6	6.9	12.2	18.0
Methamphetamine	15.7	35.4	28.8	14.2	10.1	28.5
PCP	0.0	0.0	0.0	0.0	0.0	0.0
Any Drug ^b	63.8	77.7	87.5	63.7	83.9	78.6
Any Drug (excluding alcohol)	60.1	75.0	81.3	57.6	74.4	77.2
Multiple Drugs ^c	22.0	35.4	36.5	22.3	31.4	37.6
Multiple Drugs (excluding alcohol)	18.6	33.5	34.8	18.1	29.7	34.3

ADAM is a program of the National Institute of Justice, the research, development, and evaluation agency of the U.S. Department of Justice.

^a The period of sensitivity of a urine drug test varies. Sensitivity to recent alcohol use is measured in hours; marijuana, in weeks; and cocaine, opiates, methamphetamine and PCP, in days.

b "Any drug" is defined as one or more of the six drugs listed in the table: alcohol, cocaine, marijuana, opiates, methamphetamine, or PCP.

^c "Multiple drugs" is defined as any two or more of the drugs listed in the table.

Weighted Estimates Based on Self-Report

5. Education: Highest Deg	ree (%)
None	25.1
High school or GED	42.2
Vocational or trade school	5.0
Some college or two-year associate	24.0
Four-year degree	3.7

9. Drug Use in Past 12 Months and

6. Main Residence in Past 30 Days (%)						
House, mobile home, apartment	71.5					
Group quarters	3.2					
Hospital or care facility	0.9					
Incarceration facility	2.3					
Shelter	2.8					
No fixed residence	18.6					
Other	0.8					

7. Current Employment Stat	us (%)
Working full time/ active military status	27.4
Working part time/ seasonal	13.9
Unemployed, looking for work	37.7
Unemployed, not looking for work	9.5
In school only	2.8
Retired	0.2
Disabled for work	8.5

8. Current Health Insurance	: (%)
No Insurance	60.9
Individually purchased	3.2
Employer- or union-funded	10.3
State-government funded	20.7
Retirement Medicare	0.0
Disability Medicare	3.6
Multiple types	1.3

11. Past-Year Drug Injection (%)

25.9

50.4

23.7

No drug use in past year

Used drugs, did not inject

Injected drugs

30 Days						
9a. Used in Past 12 Months	s (%)					
Alcohol	56.2					
Crack Cocaine	24.4					
Powder Cocaine	20.4					
Marijuana	60.0					
Heroin	15.8					
Methamphetamine	33.7					
9b. For 9a, Mean Number o	of Days Used in					
Past 12 Months						
Alcohol	111.0					
Crack Cocaine	76.6					
Powder Cocaine	51.1					
Marijuana	91.8					
Heroin	118.9					
Methamphetamine						
	91.5					
,	91.5					
9c. Used in Past 30 Days (

									cocaine
									marijuana
									opiates
		1							meth
100%	% 10	809)%	60	0%	40)%	20	09
	%	809)%	60	0%	40	ay Use	20 ed Seven Da ed Three Day	■ Self Repo

12. Prior Arrests in F	Past Year (%)
None	36.7
1-2	46.9
3-5	13.3
6 or more	3.1

NOTE: Comparability for opiate items is not exact. Self-reported opiate use refers specifically to heroin, while urinalysis results may include a small percentage of other opiate users.



Crack Cocaine

Marijuana

Heroin

Powder Cocaine

Methamphetamine

19.8

14.5

51.1

12.8


26.3

Weighted Estimates Based on Self-Report

13. Experience with Treatment for Drugs (including Alcohol) or Mental Health Problems													
		Inpatient Drug/Alcohol			Outpa	atient Drug/A	cohol	Inpatient Mental Health					
	Any Type of Treatment, Percent Ever		,	Mean Months In Past Year**	Percent Ever	Percent of Months in Past Year*	Mean Months In Past Year**	Percent Ever	Percent of Days in Past Year*	Mean Months In Past Year**			
Alcohol Users	70.3	49.6	8.0	2.0	42.7	38.3	5.0	16.4	3.1	2.0			
Crack Cocaine Users	86.4	74.1	10.3	2.0	50.2	42.4	5.0	20.4	3.8	2.0			
Powder Cocaine Users	86.0	65.0	12.0	2.0	51.3	29.6	4.0	25.3	1.1	1.0			
Marijuana Users	68.0	46.1	11.3	2.0	42.7	41.8	5.0	15.9	2.3	1.0			
Methamphetamine Users	81.1	59.1	15.3	3.0	55.5	44.2	5.0	21.1	6.4	2.0			
Heroin Users	85.9	74.5	11.8	2.0	55.0	31.2	4.0	26.6	2.1	1.0			

^{*} Based on those who reported receiving past-year treatment of the type indicated.

14. Percent at Risk of Abuse or Dependence



Dependence: A pattern of alcohol or drug use resulting in a combination of three or more indicators, including drinking/using more than intended, neglecting responsibilities, wanting to cut down, others object to use, frequently thinking about drinking/using, and drinking/using to relieve negative emotions; provided that thinking about drinking/using or negative emotions is included in the three or more indicators. If both thinking about drinking/using and negative emotions are the only two indicators, this also indicates dependence.

Abuse: Pattern of alcohol or drug use resulting in a combination of two indicators from those listed above; *except* if the two indicators are both thinking about drinking/using drugs and negative emotions. Also, abuse is present if three or more indicators are present but neither thinking about use nor negative emotions is reported.

Note: For both dependence and abuse, positive responses to using more than intended *and* neglecting responsibilities represent only one indicator. Further, designations are mutually exclusive. These definitions are partly based on American Psychiatric Association DSM-IV criteria.




^{**} Months with any treatment experience for those receiving past-year treatment of the type indicated.

Weighted Estimates Based on Self-Report

15. Location of Last D	rug Buy wit	h Cash in Pa	st 30 Days (%	6)	
	n	Public Building	House/ Apartment	Outdoor Area	Other Area
Crack Cocaine	70	11.9	31.7	51.4	5.0
Powder Cocaine	58	13.3	29.4	54.8	2.5
Marijuana	105	14.2	48.8	35.8	1.2
Heroin	56	5.3	20.9	72.3	1.6
Methamphetamine	88	11.7	56.2	31.1	1.1

17. Method of Ol	otainin	g Drug in Pa	ast 30 Days, r	= 353		
crack cocaine	•			Ì		
powder cocaine	: _					
marijuana						
heroir						
methamphetamine						
	0%	20%	40%	60%	80%	100%
■ cash only	■ non-c	ash only	□ cash and sor	nething else		





Dealers were

not available 34%



*Credit, front, manufacture, transport, steal, gift, other.







15%

Police activity

11%

PORTLAND — MULTNOMAH COUNTY

Adult Women: An Unweighted Sample of Booked Arrestees



Facilities Sampled: 1
Total Facilities: 1

Completed Interviews in Sample: 104 Interview Refusal Rate: 13.3% Completed Urine Tests in Sample: 92 Urine Refusal Rate: 4.8%

1. Age					
	<21	21–25	26–30	31–35	36+
	6.7%	19.2%	12.5%	19.2%	42.3%

2. Race/Ethnicity						
Black	White	Hispanic	Other			
16.8%	60.4%	12.9%	9.9%			

3. Percent with Drug-Positive Urin	ne Test ^a	by Age ar	nd Race/	Ethnicity							
	Total Sample				Age			Race/Ethnicity			
	%	95% CI*	<21	21-25	26-30	31-35	36+	Black	White	Hispanic	Other
Alcohol	5.5	N/A	0.0	0.0	9.1	0.0	10.5	0.0	5.6	16.7	0.0
Cocaine	39.6	N/A	0.0	33.3	45.5	50.0	42.1	75.0	25.9	58.3	33.3
Marijuana	35.2	N/A	50.0	50.0	0.0	33.3	36.8	25.0	40.7	25.0	33.3
Opiates	22.0	N/A	50.0	16.7	9.1	16.7	26.3	12.5	25.9	33.3	0.0
Methamphetamine	29.7	N/A	50.0	38.9	36.4	33.3	18.4	0.0	40.7	33.3	11.1
PCP	0.0	N/A	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Any Drug ^b	83.5	N/A	100.0	88.9	72.7	88.9	78.9	81.3	81.5	100.0	77.8
Any Drug (excluding alcohol)	81.3	N/A	100.0	88.9	63.6	88.9	76.3	81.3	77.8	100.0	77.8
Multiple Drugs ^c	37.4	N/A	33.3	33.3	18.2	38.9	44.7	31.3	46.3	33.3	0.0
Multiple Drugs (excluding alcohol)	36.3	N/A	33.3	33.3	18.2	38.9	42.1	31.3	44.4	33.3	0.0

4. Percent with Drug-Positive Urine Test ^a by Offense Category						
	Violent	Property	Drug	Domestic Violence	Driving While Intoxicated	Other
Alcohol	7.4	5.0	0.0	5.9	0.0	5.1
Cocaine	25.9	20.0	42.9	17.6	0.0	53.8
Marijuana	48.1	45.0	42.9	35.3	50.0	28.2
Opiates	11.1	20.0	38.1	17.6	50.0	23.1
Methamphetamine	18.5	45.0	42.9	11.8	0.0	28.2
PCP	0.0	0.0	0.0	0.0	0.0	0.0
Any Drug ^b	77.8	80.0	100.0	64.7	100.0	84.6
Any Drug (excluding alcohol)	74.1	75.0	100.0	58.8	100.0	84.6
Multiple Drugs ^c	33.3	40.0	47.6	23.5	0.0	38.5
Multiple Drugs (excluding alcohol)	29.6	40.0	47.6	23.5	0.0	38.5

ADAM is a program of the National Institute of Justice, the research, development, and evaluation agency of the U.S. Department of Justice.

^{*}Confidence intervals are not reported because female samples are not probability-based.

^a The period of sensitivity of a urine drug test varies. Sensitivity to recent alcohol use is measured in hours; marijuana, in weeks; and cocaine, opiates, methamphetamine and PCP, in days.

^b "Any drug" is defined as one or more of the six drugs listed in the table: alcohol, cocaine, marijuana, opiates, methamphetamine, or PCP.

^c "Multiple drugs" is defined as any two or more of the drugs listed in the table.

Percentages and Numbers Based on Self-Report

5. Education: Highest Deg	ree (%)
None	22.8
High school or GED	32.7
Vocational or trade school	4.0
Some college or two-year associate	36.6
Four-year degree	4.0

6. Main Residence in Past (%)	30 Days
House, mobile home, apartment	63.4
Group quarters	10.9
Hospital or care facility	4.0
Incarceration facility	1.0
Shelter	2.0
No fixed residence	18.8
Other	0.0

7. Current Employment Stat	us (%)
Working full time/ active military status	22.5
Working part time/ seasonal	3.4
Unemployed, looking for work	41.6
Unemployed, not looking for work	20.2
In school only	1.1
Retired	0.0
Disabled for work	11.2

8. Current Health Insurance	(%)
No Insurance	49.0
Individually purchased	2.0
Employer- or union-funded	9.0
State-government funded	34.0
Retirement Medicare	0.0
Disability Medicare	5.0
Multiple types	1.0

11. Past-Year Drug Injection (%)

24.0

44.8

31.3

No drug use in past year

Used drugs, did not inject

Injected drugs

 Drug Use in Past 12 M Days 	onths and
9a. Used in Past 12 Months	(%)
Alcohol	47.5
Crack Cocaine	38.6
Powder Cocaine	23.8
Marijuana	49.5
Heroin	26.7
Methamphetamine	40.6
9b. For 9a, Mean Number of Past 12 Months	Days Used in
	Days Used in 90.9
Past 12 Months	
Past 12 Months Alcohol	90.9
Past 12 Months Alcohol Crack Cocaine	90.9 121.5
Past 12 Months Alcohol Crack Cocaine Powder Cocaine	90.9 121.5 47.0
Past 12 Months Alcohol Crack Cocaine Powder Cocaine Marijuana	90.9 121.5 47.0 102.9
Past 12 Months Alcohol Crack Cocaine Powder Cocaine Marijuana Heroin	90.9 121.5 47.0 102.9 96.9
Past 12 Months Alcohol Crack Cocaine Powder Cocaine Marijuana Heroin	90.9 121.5 47.0 102.9 96.9 99.3
Past 12 Months Alcohol Crack Cocaine Powder Cocaine Marijuana Heroin Methamphetamine	90.9 121.5 47.0 102.9 96.9 99.3
Past 12 Months Alcohol Crack Cocaine Powder Cocaine Marijuana Heroin Methamphetamine 9c. Used in Past 30 Days (%	90.9 121.5 47.0 102.9 96.9 99.3

cocaine						
_						
marijuana						
-						
opiates						
meth						
0%	20	% 40	0% 60	0% 8	0% 10	00%

12. Prior Arrests in Past Year (%)						
None	40.6					
1-2	38.5					
3-5	14.6					
6 or more	6.3					

NOTE: Comparability for opiate items is not exact. Self-reported opiate use refers specifically to heroin, while urinalysis results may include a small percentage of other opiate users.



Methamphetamine

Marijuana

Heroin

40.6

17.0


30.5

Percentages and Numbers Based on Self-Report

13. Experience with Treatment for Drugs (including Alcohol) or Mental Health Problems										
		Inpatient Drug/Alcohol			Outpatient Drug/Alcohol			Inpatient Mental Health		
	Any Type of Treatment, Percent Ever	Percent Ever	Percent of Days in Past Year*	Mean Months In Past Year**	Percent Ever	Percent of Months in Past Year*	Mean Months In Past Year**		Percent of Days in Past Year*	Mean Months In Past Year**
Alcohol Users	62.5	47.9	7.2	2.0	37.5	46.7	6.0	25.0	0.7	5.0
Crack Cocaine Users	82.1	74.4	16.5	3.0	59.0	42.7	5.0	30.8	1.5	4.0
Powder Cocaine Users	79.2	79.2	10.7	2.0	37.5	27.8	3.0	29.2	1.5	1.0
Marijuana Users	62.0	46.0	7.7	2.0	32.0	75.0	9.0	22.0	1.1	4.0
Methamphetamine Users	73.2	61.0	7.4	2.0	43.9	56.3	7.0	29.3	1.3	3.0
Heroin Users	74.1	63.0	15.5	3.0	51.9	45.8	6.0	29.6	0.8	1.0

^{*} Based on those who reported receiving past-year treatment of the type indicated.

14. Percent at Risk of Abuse or Dependence



Dependence: A pattern of alcohol or drug use resulting in a combination of three or more indicators, including drinking/using more than intended, neglecting responsibilities, wanting to cut down, others object to use, frequently thinking about drinking/using, and drinking/using to relieve negative emotions; provided that thinking about drinking/using or negative emotions is included in the three or more indicators. If both thinking about drinking/using and negative emotions are the only two indicators, this also indicates dependence.

Abuse: Pattern of alcohol or drug use resulting in a combination of two indicators from those listed above; except if the two indicators are both thinking about drinking/using drugs and negative emotions. Also, abuse is present if three or more indicators are present but neither thinking about use nor negative emotions is reported.


Note: For both dependence and abuse, positive responses to using more than intended and neglecting responsibilities represent only one indicator. Further, designations are mutually exclusive. These definitions are partly based on American Psychiatric Association DSM-IV criteria.




^{**} Months with any treatment experience for those receiving past-year treatment of the type indicated.

Percentages and Numbers Based on Self-Report

15. Location of Last Drug Buy with Cash in Past 30 Days (%)										
	n	Public Building	House/ Apartment	Outdoor Area	Other Area					
Crack Cocaine	24	8.3	29.2	58.3	4.2					
Powder Cocaine	7	0.0	42.9	57.1	0.0					
Marijuana	15	6.7	53.3	26.7	13.3					
Heroin	13	7.7	23.1	69.2	0.0					
Methamphetamine	18	0.0	72.2	11.1	16.7					





*Credit, front, manufacture, transport, steal, gift, other.

