



2025 Patient Treatment Protocols

Effective January 01, 2025



Multnomah County Health Department
EMERGENCY MEDICAL SERVICES



MEMORANDUM

TO: All Multnomah County EMS Agencies

FROM: Jon Jui, MD, MPH
EMS Medical Director
Multnomah County EMS

DATE: January 01, 2025

SUBJECT: Scope of Practice
Treatment Protocol use


As EMS Medical Director for Multnomah County, I am responsible for the protocols used by Emergency Medical Services Providers (EMS Providers) working for licensed ambulance and/or 9-1-1 public safety agencies in Multnomah County. In this respect, out-of-hospital care providers must use the attached Multnomah County EMS Patient Treatment Protocols while treating patients in Multnomah County.

The protocols are defined as the Preface, Patient Treatment, Drugs, Procedures, Communications, Operations, Trauma and Special Operations sections.

EMS Providers may function up to but not outside of the full scope of practice for the level at which they are licensed as defined in the Oregon Administrative Rules (OAR 847-035-0030).

The scope of practice for Multnomah County EMS Protocols is indicated in Multnomah County Patient Treatment Protocol 00.010 (EMS Provider Scope of Practice). However, EMS Providers are not covered for activities outside of the scope of their employment unless otherwise specifically authorized.

Effective Date: January 01, 2025



Jon Jui, MD, MPH
EMS Medical Director
Multnomah County

619 NW 6th Avenue
Portland, OR 97209
(503) 988-3220 phone
(503) 988-4017 fax

<https://multco.us/health/providers/emergency-medical-services>

CONTENTS

Treatment

(Section 10.000)

Drugs

(Section 20.000)

Procedures

(Section 30.000)

Communications

(Section 40.000)

Operations

(Section 50.000)

Trauma Protocol EMSAB

(Section 60.000)

Special Operations

(Section 70.000)

Table of Contents

Preface Section	00.000
EMS Provider Scope of Practice and Nomenclature	00.010
Death in the Field	00.020
Death Notification	00.025
Dying and Death, POLST, Do Not Attempt Resuscitation Orders	00.030
Medical Control for Drugs and Procedures	00.040
Treatment	Section 10.000
Abdominal Pain	10.010
Altered Mental Status and Coma.....	10.020
Anaphylaxis and Allergic Reaction.....	10.030
Burns	10.040
Cardiac Arrest	10.050
Emergency Medical Responder/EMT	
Paramedic/EMT-Intermediate	
Quick Reference to Pediatric Drugs	
Cardiac Dysrhythmias.....	10.060
Chest Pain/Acute Coronary Syndrome.....	10.070
Crush Injury	10.075
Diabetic Emergencies	10.076
Epistaxis.....	10.077
Eye Emergencies.....	10.078
Hyperthermia.....	10.080
Hypothermia.....	10.090
Musculoskeletal Injuries	10.100
Nausea and Vomiting	10.110
Neonatal Resuscitation.....	10.120
OB/GYN Emergencies.....	10.130
Pain Management	10.135
Poisons and Overdoses	10.140
Psychiatric and Behavior Disorders.....	10.150
Respiratory Distress.....	10.160
Seizures	10.170
Sepsis	10.175
Shock.....	10.180
Stroke/CVA.....	10.190
Submerged Patient	10.200
Trauma Patient Evaluation and Treatment.....	10.300
Drugs	Section 20.000
Acetaminophen.....	20.005
Activated Charcoal	20.010
Adenosine (Adenocard®)	20.020

Drugs (cont.)	Section 20.000
Albuterol (Ventolin®)	20.030
Alprazolam (Xanax)	20.035
Amiodarone	20.040
Aspirin	20.050
Atropine Sulfate	20.060
Buprenorphine	20.065
Calcium Gluconate	20.070
Dexamethasone (Decadron®)	20.075
Dextrose 50% / 10%	20.080
Diphenhydramine (Benadryl®)	20.090
Epinephrine	20.110
Esmolol (Brevibloc®)	20.112
Etomidate (Amidate®)	20.115
Fentanyl (Sublimaze)	20.117
Furosemide (Lasix®)	20.120
Glucagon	20.130
Glucose, Oral	20.140
Haloperidol (Haldol®)	20.142
Hydroxocobalamin (Cyanokit®)	20.145
Ibuprofen	20.147
Ipratropium Bromide (Atrovent®)	20.150
Ketamine	20.155
Ketorolac (Toradol®)	20.157
Labetalol	20.158
Lidocaine (Xylocaine®)	20.160
Magnesium Sulfate	20.170
Midazolam (Versed®)	20.190
Morphine Sulfate	20.200
Naloxone (Narcan®)	20.210
Nitroglycerin	20.220
Norepinephrine (Levophed®)	20.225
Olanzapine (Zyprexa®)	20.227
Ondansetron (Zofran®)	20.230
Oxygen	20.240
Oxymetazoline hydrochloride (Afrin®)	20.245
Oxytocin	20.247
Pralidoxime (Protopam/2-PAM)	20.250
Prochlorperazine (Compazine)	20.253
Proparacaine	20.254
Rocuronium Bromide	20.255
Sodium Bicarbonate	20.260
Sodium Thiosulfate	20.265
Succinylcholine	20.270
Sufentanil (Sufenta)	20.275

Tranexamic Acid (TXA)	20.277
Vecuronium Bromide (Norcuron®)	20.290
Whole Blood (LTOWB+)	20.295
Xylocaine, Viscous (Lidocaine®, topical)	20.300
Ziprasidone (Geodon)	20.310
Procedures	Section 30.000
Airway Management	30.010
Automatic Implantable Cardio-Defibrillator (AICD) Deactivation	30.020
Continuous Positive Airway Pressure (CPAP)	30.032
Endotracheal Intubation.....	30.040
Emergency Surgical Cricothyrotomy	30.060
Double Sequential External Defibrillation	30.065
End-Tidal CO ₂ Monitoring	30.070
i-gel Supraglottic Airway Device.....	30.075
Induced Hypothermia	30.077
Intraosseous Infusion.....	30.080
Intravenous Lines and IV Solutions Management	30.090
Intravenous Solutions Control and Monitoring.....	30.100
Left Ventricular Assist Device Troubleshooting (HVAD & LVAD)	30.107
LUCAS Chest Compression Device	30.109
Patient Disposition.....	30.110
Patient Disposition Flow Chart	
Discharge from Scene form	
Pediatric Field Initial Survey.....	30.130
Pelvic Wrap	30.132
Physical Restraint	30.136
PICC Line Access.....	30.140
Positive End Expiratory Pressure (PEEP)	30.145
Selective Spinal Immobilization.....	30.150
Sports Equipment Removal	30.160
Removal of Taser Barbs.....	30.162
Tension Pneumothorax Decompression (Thoracentesis)	30.170
Transcutaneous Pacing	30.180
Unity Center for Behavioral Health Emergencies	30.190
Video Laryngoscope.....	30.200
Whole Blood (LTOWB+)	30.210
Communications	Section 40.000
BOEC EMS Dispatch (Multnomah County)	40.010
Medical Resource Hospital, Receiving Hospitals and the Trauma Communications Center	40.020
Operations.....	Section 50.000
Cancellation/Slow Down Policy	50.010

Operations (cont.).....	Section 50.000
Crime Scene Response.....	50.020
Diversion System.....	50.030
Documentation of Care.....	50.040
EMS Management of Lower Acuity Calls	50.045
Field Surgical Response Team	50.050
Hazardous Materials	50.060
Multiple Toxic Exposure	50.070
Helicopter Ambulance Service (HEMS).....	50.080
Mass Casualty Incident (MCI)	50.100
Task Card – Medical	
Scene Checklist	
Task Card – Treatment	
Treatment Area Guidelines	
Scene Checklist	
Task Card – Triage	
Task Card – Transportation	
Task Card – Destination	
Task Card – Regional Hospital Coordinator	
Patient Transport Log	
On-Scene Medical Control.....	50.110
Staging for High Risk Response	50.120
Transport by Fire Apparatus.....	50.130
Transportation and Patient Destination.....	50.140
Trauma Protocol EMSAB	Section 60.000
Trauma System Patient Entry Criteria	60.010
Medical Direction	60.020
Communications.....	60.030
Transport Protocol	60.040
Mode of Transport	60.050
Recommended Air Ambulance Response Zones.....	60.060
Patient Evaluation Protocol	60.070
Scene Time	60.090
EMSAB Designated Hospitals, Trauma Line Map	60.100
Special Operations.....	Section 70.000
Special Operations Treatment	
Buprenorphine (Suboxone)	70.010
Hydrogen Cyanide (HCN)	70.020
Hydrogen Fluoride	70.030
Organophosphates.....	70.040
Special Operations Procedures	
Field Amputation	70.045
Ventilator Management.....	70.047
Impact Uni-Vent Model 73X.....	70.050
XSTAT	70.090

PREFACE

EMS Provider Scope of Practice

The Oregon Medical Board is responsible for the scope of practice of Emergency Medical Service Providers as well as the requirements and duties of these providers' supervising physicians (EMS Medical Directors), to include an ambulance-based clinician (EMT, paramedic, nurse, nurse practitioner, physician, physician assistant).

To view the complete Scope of Practice, see Oregon Administrative Rule (OAR) 847-035-0030.

Scope of Practice: Each EMS Provider can perform the procedures of each lower level certification plus those listed in their certification level.

Airway and Breathing

Emergency Medical Responder	Emergency Medical Technician	Advanced EMT	EMT-Intermediate	Paramedic or ambulance based clinician
<ul style="list-style-type: none"> BVM NPA OPA Pharyngeal suctioning 	<ul style="list-style-type: none"> Supraglottic airway device Tracheo-bronchial suctioning of ET tube Ventilator 			<ul style="list-style-type: none"> Cricothyrotomy Endotracheal intubation

Pharmacological Intervention

Emergency Medical Responder	Emergency Medical Technician	Advanced EMT	EMT-Intermediate	Paramedic or ambulance based clinician
<ul style="list-style-type: none"> Aspirin for cardiac Epi by auto-injector for anaphylaxis Naloxone via IN device or auto-injector Oral Glucose Oxygen 	<ul style="list-style-type: none"> Activated charcoal Epi by IM injection for anaphylaxis Nebulized albuterol, levalbuterol and/or DuoNeb Nitroglycerine (assist patient with own NTG) Pralidoxime and Atropine by autoinjector for organophosphate agents 	<ul style="list-style-type: none"> Dextrose, hypertonic Glucagon IO lidocaine (anesthetic) Immunizations Ipratropium Isotonic IV solutions Naloxone Nitroglycerine Nitrous oxide Tuberculosis skin testing 	<ul style="list-style-type: none"> Amiodarone Atropine Benzodiazepines for seizure or agitation Diphenhydramine Epinephrine Fentanyl Furosemide Lidocaine Morphine Zofran 	<ul style="list-style-type: none"> Administer medications authorized by protocol or physician's direct order

Medical/Cardiac Care

Emergency Medical Responder	Emergency Medical Technician	Advanced EMT	EMT-Intermediate	Paramedic or ambulance based clinician
<ul style="list-style-type: none"> • AED / SAED • Hemorrhage Control 	<ul style="list-style-type: none"> • Blood glucose monitoring 	<ul style="list-style-type: none"> • Blood draws • Start IO's in pediatrics and adults • Start IVs or saline locks 	<ul style="list-style-type: none"> • EKG interpretation • Manual defibrillation • Orogastric tube 	<ul style="list-style-type: none"> • Access indwelling catheters and implanted central IV ports • Cardioversion • Nasogastric tube • Needle Thoracostomy • Transcutaneous Pacing

EMS Provider Nomenclature

The term “EMS Provider” has replaced the term “EMT” to describe all scope of practice levels. The term “EMT” now refers to the level formerly called “EMT-Basic.” Per Oregon Administrative Rules 333-265-0000 (Definitions) and Oregon Revised Statute Chapter 682 the terms have changed as follows:

Old	New
Did not exist	Advanced Emergency Medical Technician (AEMT or Advanced EMT) • Is not used in these protocols
EMT	Emergency Medical Services Provider or EMS Provider
EMT-B	EMT or Emergency Medical Technician
EMT-I	Remains the same (or EMT-Intermediate)
EMT-P	Paramedic
First Responder	Emergency Medical Responder or EMR

Death in the Field

Purpose:

To define under what conditions treatment can be withheld or stopped.

A. Resuscitation efforts may be withheld if:

1. The patient has a “DNAR” order.
2. The patient is pulseless and apneic in a mass casualty incident or multiple patient scene where the resources of the system are required for the stabilization of living patients.
3. Decapitation.
4. Rigor mortis in a warm environment.
5. Decomposition.
6. Skin discoloration in dependent body parts (dependent lividity).

B. Traumatic Cardiac Arrest

1. A victim of trauma (blunt or penetrating) who has no vital signs at the scene may be declared dead. If opening the airway does not restore vital signs/signs of life, the patient should NOT be transported unless there are extenuating circumstances.
2. A cardiac monitor may be beneficial in determining death in the field when you suspect a medical cause or hypovolemia:
 - a. An organized rhythm may suggest profound hypovolemia and/or tension pneumothorax, and may respond to fluid resuscitation and/or needle decompression, respectively.
 - b. VF should raise your index of suspicion for a medical event.
3. At a trauma scene consider the circumstances surrounding the incident, including the possibility of cardiac arrhythmia, seizure, or hypoglycemia. When a medical event is suspected, treat as a medical event.
4. If the patient deteriorates to no vital signs (i.e., no pulse/respiration), a cardiac monitor should be applied. A viable rhythm especially in patients with penetrating trauma may reflect hypovolemia or obstructive shock (tamponade, tension pneumothorax) and aggressive treatment should be continued.
5. If a patient deteriorates to cardiac arrest during transport, perform CPR and notify the trauma facility.

C. Medical Cardiac Arrest

1. If the initial EKG shows asystole or agonal rhythm confirmed in 6 leads with full gain, and the patient in the responder's best judgment would not benefit from resuscitation:
 - a. The PIC may determine death in the field; OR
 - b. Begin BLS procedures, and contact OLMC with available patient history, current condition, and with a request for advice regarding discontinuing resuscitation.
2. The PIC may determine the patient to be dead in the field if the patient persists in asystole (confirmed in 6 leads with full gain) after the airway is established, and the asystole protocol has been exhausted.
3. Death in the field may be determined with EtCO₂ of < 10 in patients with PEA after 20 minutes of ACLS resuscitation. For patients with EtCO₂ > 10 either continue resuscitation or contact OLMC to stop resuscitation.
4. Patients in VF should be treated and transported.

Notes and Precautions

- A. ORS allows a layperson, EMT or paramedic to determine "Death in the Field."
- B. Consult OLMC with any doubt about the resuscitation potential of the patient.
- C. A person who was pulseless or apneic and has received CPR and has been resuscitated is not precluded from later being a candidate for solid organ donation.

Death Notification

Purpose:

To establish guidelines for the handling of the body and required notification following a declaration of death as outlined in ORS Chapter 146. The goal of an investigation by the medical examiner's office is to determine the cause and manner of death.

Procedures:

- A. If the patient appears to meet obvious death in the field criteria, have only one person enter the scene to verify death; limit access if possible. Don't move the decedent unless necessary. Document anything that was altered by your examination (e.g. unbuttoned/removed clothing, movement of the decedent, etc.).
- B. Contact police for all deaths in the field except for hospice and skilled nursing facilities.
- C. Upon declaration of death, the medical examiner (ME) must be contacted. Until contact is made with the ME:
 - 1. Do not move the body.
 - 2. Do not cover the body unless necessary (outside, public place). If covering the body is necessary, use a new/clean non-cloth disposable sheet or blanket such as an emergency blanket.
 - 3. Do not remove clothing or cleanse the body or otherwise alter the appearance of the state of the body.
 - 4. Do not remove any of the effects of the deceased or instruments or weapons related to the death.
 - 5. Do not let anyone in the area where the deceased is located.
 - 6. If resuscitation was attempted, do not remove IV's, advanced airways, or defib/ECG pads. Circle all IV attempts or any trauma or marks that you caused to the body with an ink pen if possible.
- D. Depending on the circumstances, the ME will either respond to the scene for a full investigation or release the body to a funeral home with a limited investigation. Generally, it is best to turn the scene over to law enforcement once you have given a report.
- E. You should not leave the scene without passing the scene off to law enforcement or until the ME has released you over the phone or the ME arrives at the scene and has released you.
- F. The following documentation is required for declaration of death calls:
 - 1. Location and position the body was found.
 - 2. Location of evidence if moved for safety concerns (gun, knife, bat, etc.).
 - 3. Anything suspicious (e.g. bruises on the body, deformed arm, black eye, comments made by bystanders/relatives/friends, etc.).
 - 4. Name and title of individual the scene is turned over to (law enforcement, ME, another crew) and the disposition of the body.
 - 5. The name of the ME if the body is released with a limited investigation.
 - 6. Follow your individual agency's medical records policy for listing witnesses or possible witnesses with contact information.

Notes

- A.** Once the person is declared dead, your jurisdiction ends. Even law enforcement is not allowed to touch or move the body. Only the ME, Deputy ME (also referred to as a Medicolegal Death Investigator), or District Attorney, has lawful authority over the body. Any of these individuals can grant access or removal of the body.
- B.** Not all deaths are under the jurisdiction of the ME (e.g. patient on hospice care longer than 24 hours, patient who dies in a skilled nursing facility). However, EMS calls should be considered an ME case and reported to the ME. It is best to let the ME decide if this is their case or not.
- C.** Your chart may be read by the ME's office and if read, will become part of the report for cause and manner of death.
- D.** In smaller counties and jurisdictions, law enforcement officers may be appointed as Deputy ME's or medicolegal Death Investigators, who under the direction of the ME's office, can investigate deaths and authorize the removal of a body of a deceased person from the apparent place of death.
- E.** If you suspect a COVID-19 death, document the names and contact information of everyone who had contact with the person that is on scene.
- F.** The following information should be available, if possible, prior to contacting the ME. The ME may not ask for all this information but be ready with this information.
 - Your name
 - Unit number
 - What you were dispatched on
 - How you found the patient
 - Brief description of your actions
 - Whether you suspect foul play
 - Whether death occurred at work
 - Whether death occurred while in custody
 - Whether death was the result of a crime
 - Whether death was unattended
 - Whether cause of death might be from a contagious disease
 - Any evidence of drug use
 - Name of deceased
 - Address of deceased
 - Age of deceased
 - Gender of deceased
 - Medical history
 - Medications
 - Primary caregiver and phone number
 - Family contact
 - Funeral home

Dying and Death, POLST, and Do Not Attempt Resuscitation Orders

A. POLST ORDERS AND DECISION MAKING

1. In the pulseless and apneic patient who does not meet DEATH IN THE FIELD criteria, but is suspected to be a candidate for withholding resuscitation, begin CPR and contact OLMC.
2. A patient with decision-making capacity or the legally authorized representative has the right to direct his or her own medical care and can change or rescind previous directives.
3. EMS providers may honor a Do Not Attempt Resuscitation (DNAR) order signed by a physician, naturopathic physician, nurse practitioner, or physician assistant. DNAR orders apply only to the patient in cardiopulmonary arrest and do not indicate the types of treatment that a person not in arrest should receive. POLST was developed to convey orders in other circumstances.
4. Physician Orders for Life-Sustaining Treatment (POLST):
 - a. The POLST was developed to document and communicate patient treatment preferences across treatment settings. While these forms are most often used to limit care, they may also indicate that the patient wants everything medically appropriate done.
Read the form carefully!
 - b. When signed by a physician (MD or DO), naturopathic physician, nurse practitioner, or physician assistant, POLST is a medical order and EMS providers are directed to honor it in their Scope of Practice unless they have reason to doubt the validity of the orders or the patient with decision-making capacity requests change. If there are questions regarding the validity or enforceability of the health care instruction, begin BLS treatment and contact OLMC [OAR 847-035-030 (7)].
 - c. If the POLST is not immediately available, a POLST form as documented in the Electronic POLST registry hosted at MRH (503-494-7333) may also be honored.
 - **Section A:** Applies only when patient is in cardiopulmonary arrest
 - **Section B:** Applies in all other circumstances
 - For a POLST form to be valid it must include:
 - i. Patient's name
 - ii. Date signed (forms do not expire)
 - iii. Health care professional's signature (patient signature is optional)
5. The legally authorized representative may make decisions for the patient who is unable to make medical decisions. However, when in doubt or for unresolved conflict on the scene contact OLMC. The order is:
 - a. A legal guardian
 - b. A power of attorney for health care as designated by the patient on the Oregon advance directive
 - c. Spouse or legal domestic partner
 - d. Adult children
 - e. Parent

Dying and Death, POLST, and Do Not Attempt Resuscitation Orders

6. Death with Dignity

If a person who is terminally ill and appears to have ingested medication under the provisions of the Oregon Death with Dignity Act, the EMS provider should:

- a. Provide comfort care as indicated.
- b. Determine who called 9-1-1 and why (i.e., to control symptoms or because the person no longer wishes to end their life with medications).
- c. Establish the presence of DNAR orders and/or documentation that this was an action under the provisions of the Death with Dignity Act.
- d. Contact OLMC.
- e. Withhold resuscitation if: DNAR orders are present, and there is evidence that this is within the provisions of the Death with Dignity Act and OLMC agrees.

B. PATIENTS ENROLLED IN HOSPICE AND DYING PATIENTS

1. Look for POLST forms (contact Registry if needed) and attempt to honor patient preferences. Always provide comfort measures.
2. If patient is enrolled in hospice and the patient has not already done so, contact hospice if possible.
3. EMS providers cannot take medical orders from a hospice nurse but their advice is often invaluable and may be followed with direction from OLMC.
4. Treat dying persons with warmth and understanding. Do not avoid them. Allow them to discuss their situation, but do not push them to talk.
5. Many dying people are not upset by discussions of death as long as you do not take away all of their hope.
6. Touching a dying person is important. Use words like “death.” Do not use meaningless synonyms.
7. Ask the person how you might help.
8. Give factual information.
9. Be aware of your own fears regarding death and admit when a dying person reminds you of a loved one. If a particular person is too disturbing, have other members of the responding team take over.
10. Consider providing pain/symptom management and not transporting patient if they are Comfort Measures Only, the symptoms can be managed, and the patient and caregivers on scene do not want transport to the hospital. Consider OLMC contact for advice.

C. CARE OF GRIEVING PERSONS

Resuscitation phase

1. As time allows give accurate and truthful updates about the patient’s prognosis. If available, assign one person to interact with and support family members.
2. Consider gently removing children from the resuscitation area.
3. Depending upon the emotional state of family members, consider allowing them to watch and/or participate in a limited and appropriate way.

Dying and Death, POLST, and Do Not Attempt Resuscitation Orders

4. If family or friends were doing CPR prior to your arrival, commend their efforts.
5. If family or friends are disruptive consider removing them or try assigning simple tasks, such as helping bring in the stretcher, telling other family about the event, and calling the doctor or minister.
6. Be respectful. Make requests. Don't give orders.

Once death is determined

- a. Treat the recently dead with respect.
- b. Tell family and friends of the death honestly. Use the words "death" or "dead."
Avoid using euphemisms such as "passed away" or "gone."
- c. Avoid using past tense terms when speaking to survivors of the recently dead.
- d. Allow family and friends to express their emotions. Listen to them if they want to talk but don't push them.
- e. Give factual information.
- f. Genuine warmth and compassion will be more helpful than almost anything else for survivors. Don't feel it necessary to say the "right" things. Listening often provides grieving people with the most comfort.

Focusing on survivors

- a. See to it that survivors have a support system present before you leave. Consider calling TIP through EMS Dispatch. Call friends, family, clergy, or neighbors to be with them. Respect the survivors' wishes to be alone.
- b. Explain the next steps to them after you have pronounced death. This will include the police coming to make reports, possibly the medical examiner, and the possible need for an autopsy.
- c. Before moving or altering the body, contact the Medical Examiner's office as soon as possible.
- d. Allow family and friends to say their good-byes if possible.
- e. A chaplain may be helpful in assisting with survivors. Call early, as chaplains do not have code-3 capabilities.
- f. Help survivors make decisions such as which people should be called. If they ask you to make calls, try to comply. Mention the need to find a funeral home, if one has not been chosen. Clergy may also be helpful with this decision.

C. DEATH OF A CHILD

1. Do not accuse the parents of abuse or neglect, but note carefully the patient's surroundings and the general physical condition of the child.
2. Do not be overly silent, which may imply guilt to the parents.
3. Ask the parents only necessary questions and do not judge or evaluate them. Do not tell them what they "should have" been doing before your arrival.
4. Remind parents to arrange for child care of other children.
5. Listen carefully to their statements and answer with accurate information.
6. If there is a police investigation, tell the parents that this is routine.
7. Successful management of child deaths requires supportive, compassionate, and tactful measures.

Medical Control for Drugs and Procedures

Policy:

If a patient receives a procedure or medication; is conscious (or regains consciousness); and refuses transport, every effort shall be made to encourage transport of the patient. If the patient persists in refusing transport, see the *Non-Transport* procedure.

These protocols contain Category A and B drugs and procedures. Before using any Category B drug or procedure you **must** contact OLMC. If the EMS Provider is unable to contact OLMC, Category B drugs or procedures should be administered as indicated in the protocol. If a Category B drug or procedure is used without OLMC contact, **a written report must be sent to the Medical Director or Physician Supervisor**. Continued attempts must be made to reach OLMC en route.

Category A:

Drug or procedure will be used at the EMS Provider's discretion in accordance with the standing orders.

Drugs:

Acetaminophen	Labetalol
Activated Charcoal (ASA and APAP only)	Lidocaine (Xylocaine®)
Adenosine (Adenocard®)	Magnesium Sulfate
Albuterol	Midazolam (Versed®)
Alprazolam (Xanax)	Morphine
Amiodarone	Naloxone (Narcan®)
Aspirin	Nitroglycerin
Atropine Sulfate	Norepinephrine (Levophed®)
Calcium Gluconate	Olanzapine (Zyprexa®)
Dexamethasone (Decadron®)	Ondansetron (Zofran®)
Dextrose 10%, 50% IV	Oxymetazoline hydrochloride (Afrin)
Diphenhydramine (Benadryl®)	Oxygen
Epinephrine	Oxytocin
Esmolol (Brevibloc®)	Pralidoxime (Protopam/2-PAM)
Etomidate (Amidate®)	Prochlorperazine (Compazine)
Fentanyl (Sublimaze)	Proparacaine
Furosemide (Lasix®)	Rocuronium Bromide
Glucagon	Sodium Bicarbonate
Glucose, Oral	Sodium Thiosulfate
Haloperidol (Haldol®)	Succinylcholine
Hydroxocobalamin (Cyanokit®)	Sufentanil (Sufenta)
Ibuprofen	Tranexamic Acid (TXA)
Ipratropium (Atrovent®)	Vecuronium Bromide (Norcuron®)
IV Solutions	Whole Blood (LTOWB+)
Ketamine	Xylocaine, Viscous
Ketorolac (Toradol®)	Ziprasidone (Geodon)

Category A:**Procedures:**

Chemical Patient Restraint
Continuous Positive Airway Pressure (CPAP)
End-Tidal CO₂ Monitoring
Endotracheal Intubation
i-gel
Induced Hypothermia
Intraosseous Infusion
Intravenous Lines and IV Solutions (Management of)
Intravenous Solutions (Control and Monitoring of)
King LT-D/LTS-D Airway Device
Left Ventricular Assist Device
LUCAS
Paralytic Intubation: Advanced Airway Training Required
Physical Patient Restraint
Self-Care Instructions
Positive End Expiratory Pressure (PEEP)
Pelvic Wrap
Selective Spinal Immobilization
Sports Equipment Removal
Surgical Cricothyrotomy
Synchronous Cardioversion
 A. Unstable V-Tachycardia, OR
 B. SVT, unstable patient with BP less than 90 mm/Hg
Taser Barb Removal
Tension Pneumothorax Decompression
Transcutaneous Pacing
Video Laryngoscope

Category B:

Drug or procedure, not included in Category A, shall be initiated by request from EMS Provider to OLMC. Confirmation of dosage or procedure will be obtained directly from a Physician on Duty at OLMC.

Drugs:

Glucagon (Beta Blocker OD)
Magnesium Sulfate (OB/GYN)

Procedures:

Automatic Implantable Cardio-Defibrillator Deactivation (AICD)

Required Multnomah County Medications

- ☐ Activated Charcoal: 25 grams
- ☐ Adenosine: 6 mg, 12 mg (3 mg/mL)
- ☐ Albuterol: 2.5 mg/3 mL
- ☐ Alproazolam (Xanax) 0.25 mg tablets
- ☐ Amiodarone: 150 mg/3 mL
- ☐ Aspirin: 81 mg tablets
- ☐ Atropine: 1 mg/10 mL
- ☐ Calcium Gluconate: 10% (100 mg/mL)
- ☐ Dexamethasone: 10 mg/mL
- ☐ Dextrose:
 - ☐ 10%, 250 mL (0.1 g/mL)
 - ☐ 50%, 50 mL (0.5 g/mL)
 - ☐ Oral, 24 gram
- ☐ Diphenhydramine: 50 mg/mL
- ☐ Epinephrine:
 - ☐ 1:1,000, 1 mg/mL
 - ☐ 1:10,000, 1 mg/10 mL
- ☐ Esmolol: 100 mg/10 mL
- ☐ Etomidate: 40 mg (2 mg/mL)
- ☐ *Fentanyl: 100 mcg/2 mL
- ☐ Furosemide: 40 mg/4 mL
- ☐ Glucagon: 1 mg/mL
- ☐ Haloperidol: 5 mg/mL
- ☐ Ipratropium: 0.5 mg/2.5 mL
- ☐ *Ketamine: 500 mg/10 mL
- ☐ Ketorolac: 30 mg/3 mL
- ☐ Labetalol: 100 mg/20 mL
- ☐ Lidocaine:
 - ☐ 2%, 100 mg/5 mL
 - ☐ 2%, Viscous Jelly
- ☐ Magnesium Sulfate 50%: 1 g/2 mL
- ☐ *Midazolam: 10 mg/2 mL
- ☐ Naloxone: 2 mg/2 mL
- ☐ Nitroglycerin:
 - ☐ Tablets, 0.4 mg
 - ☐ IV Nitroglycerin 50 mg/10 mL
- ☐ Norepinephrine: by IV infusion pump only
- ☐ Olanzapine:
 - ☐ Tablets, 10 mg
- ☐ Ondansetron:
 - ☐ 4 mg/2 mL
 - ☐ Tablets, 4 mg
- ☐ Oxytocin: 10 units/mL
- ☐ Prochlorperazine (Compazine):
 - ☐ 10 mg/2 mL
- ☐ Proparacaine 0.5% solution
- ☐ Rocuronium Bromide:
 - ☐ 10mg/mL (5 mL, 10mL)
 - [9-1-1 Responding Units Only]*
- ☐ Sodium Bicarb: 50 mEq/50 mL
- ☐ Succinylcholine: 20 mg/mL
- [9-1-1 Responding Units Only]*
- ☐ *Sufentanil (Sufenta): 50 mcg/1 mL
- ☐ Tranexamic Acid (TXA): 1 gram/10 mL
- ☐ Vecuronium: 10 mg-powder (1 mg/mL)
- [9-1-1 Responding Units Only]*
- ☐ Ziprasidone: 20 mg/mL

* Must be locked and counted at each shift change

Special Operations Medications

- ☐ Pralidoxime (Protopam/2-PAM)
- ☐ Proparacaine Hydrochloride Ophthalmic Solution USP, 0.5%

NOTE:

Alternative formulations allowed with approval of EMS Medical Director.